

MINI-FOCUS ISSUE: CONGENITAL HEART DISEASE

BEGINNER

IMAGING VIGNETTE: CLINICAL VIGNETTE

# Giant Pulmonary Artery Aneurysm in Bicuspid Pulmonary Valve



## Does the Right Side Mimic the Left One?

Concetta Di Nora, MD,<sup>a</sup> Rosa Pecoraro, MD,<sup>b</sup> Arturo Evangelista, MD,<sup>c</sup> Francesco Antonini-Canterin, MD<sup>d</sup>

### ABSTRACT

Bicuspid pulmonary valve is a rare echocardiographic finding, particularly if not associated with other congenital heart diseases. We report the incidental case of a severe giant pulmonary arterial aneurysm associated to bicuspid pulmonary valve in an asymptomatic 79-year-old patient. Multimodality cardiac imaging was important for the correct diagnosis and to exclude any other potential complication. (**Level of Difficulty: Beginner.**) (J Am Coll Cardiol Case Rep 2020;2:1720-2)  
© 2020 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**A**ortic and pulmonary valves are known to stem from the same embryonic arterial trunk, and this could explain the similar presentation of some types of congenital defects in both of the 2 valves. However, due to the limited number reported, more in autoptic analysis, little is known about the effective risk and management of pulmonary abnormalities. Multimodality cardiac imaging is mandatory to better define the valve morphology and to rule out related complications. Overall, surgery remains the cornerstone of therapy, albeit definite surgical thresholds are still lacking.

### CASE REPORT

A 79-year-old patient presented to the emergency department for fever and chest pain, described as breathing and cough variable and almost continuous for weeks. No previous cardiovascular history was known, except for arterial hypertension treated with calcium antagonist therapy. A transthoracic echocardiogram performed to investigate a systolic murmur showed a moderately stenotic pulmonary valve and severe pulmonary artery aneurysm (PAA) (**Figures 1A and 1B, Videos 1 and 2**). Noninvasive estimation of pulmonary artery pressure resulted in the normal range. The chest pain was explained by a pleuritic inflammation detected at chest x-ray, and a urine test revealed an infection treated with antibiotics. A computed tomography (CT) scan was urgently performed to rule out any acute emergency, although the patient never reported any symptoms related to the PAA compression. The CT and the 3-dimensional CT reconstruction confirmed these data (**Figures 1C and 1D, Video 3**), as far as the magnetic resonance imaging (**Figure 1E**, arrow; **Videos 4, 5, 6, and 7**).

From the <sup>a</sup>Cardiothoracic Department, Azienda Sanitaria Universitaria Integrata, Udine, Italy; <sup>b</sup>Cardiology Department, AAS 5 Friuli Occidentale, Pordenone, Italy; <sup>c</sup>Department of Cardiac Imaging, Hospital Val d'Hebron, Barcelona, Spain; and the <sup>d</sup>Rehabilitative Cardiology, ORAS, Motta di Livenza, Italy. The authors have reported that they have no relationships relevant to the contents of this paper to disclose. Sarah Moharem-Elgamal, MD, PhD, served as Guest Associate Editor for this paper. The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the *JACC: Case Reports* [author instructions page](#).

Manuscript received April 25, 2020; revised manuscript received May 14, 2020, accepted May 29, 2020.


Balancing potential risks and benefits, corrective heart surgery had been proposed to the patient, but was rejected in the absence of symptoms. Currently, after 3 years of follow-up, she is asymptomatic, taking bisoprolol 5 mg/die and routinely transthoracic echocardiogram has been scheduled every 6 months.

**ABBREVIATIONS  
AND ACRONYMS**

**CT** = computed tomography scan

**PAA** = pulmonary artery aneurysm

**FIGURE 1** Patient's TTE, CT, and CMR



**(A and B)** Patient's transthoracic echocardiogram (TTE). The pulmonary trunk was severely dilated (maximum diameter of 6.39 cm), with severe dilatation of the pulmonary arteries (2.97 and 2.8 cm, respectively). The pulmonary cusps appeared myxomatous and calcific. The maximum velocity recorded was 2.98 m/s with a maximum gradient >35 mm Hg. **(C and D)** Patient's computed tomography scan (CT). The CT confirms the presence of a severe dilation of the PA trunk (6.7 × 5.9 cm), right pulmonary artery (3.8 × 3.5 cm), and left pulmonary artery (3.6 × 3.4 cm) with no sign of dissection of the wall. The pulmonary valve appeared morphologically bicuspid. The 3-dimensional CT reconstruction confirmed the severe dilatation of the pulmonary trunk and pulmonary arteries. **(E and F)** Patient's cardiac magnetic resonance imaging (CMR). The pulmonary valve appeared morphologically bicuspid, and the presence of a severe dilatation of the PA trunk and pulmonary arteries was confirmed.

## DISCUSSION

---

Bicuspid pulmonary valve is an exceedingly rare finding usually associated with PAA that could result as the consequence of chronic stenosis of the valve. Few sporadic cases of PAA have been reported until now, especially concerning aortic analysis, but such a giant PAA with bicuspid pulmonary valve in an asymptomatic woman has never been described. Moreover, as demonstrated here, a multimodality imaging approach was crucial to better define the morphology of the pulmonary valve, to exclude any other congenital defects, such as conotruncal anomalies, and to rule out any associated complications. Different cutoffs for PAA have been proposed, from a range of 26.9 mm in women (1) to 29 mm (2) in the general population, and an absolute diameter  $\geq 55$  mm has been designated for surgical intervention (3); so our case, where PAA was 64 mm, was clearly enclosed in these ranges. Overall, surgery remains the cornerstone of these pulmonary abnormalities; however, there are no clear guidelines yet on the optimal management of these patients (4). In our case, the absence of complications and symptoms has led the patient to give up the proposed corrective surgery, so we have planned a follow-up every 6 months. Moreover, albeit not clearly specified, a pharmacological treatment with beta-blockers has been introduced to reduce vascular wall stress, as is usually also done for the aneurysms of the aortic tract.

---

**ADDRESS FOR CORRESPONDENCE:** Dr. Concetta Di Nora, Cardiothoracic Department, Azienda Sanitaria Universitaria Integrata di Udine, 33010 Udine, Italy. E-mail: [concetta.dinora@gmail.com](mailto:concetta.dinora@gmail.com).

---

## REFERENCES

1. Gallego P, Rodríguez-Puras MJ, Gotarredona PS, et al. Prevalence and prognostic significance of pulmonary artery aneurysms in adults with congenital heart disease. *Int J Cardiol* 2018;270:120-5.
2. Truong QA, Massaro JM, Rogers IS, et al. Age and gender-specific pulmonary artery measurements by multi-detector computed tomography: Framingham heart study. *J Am Coll Cardiol* 2010;55:10A.
3. Kreibich M, Siepe M, Kroll J, Höhn R, Grohmann J, Beyersdorf F. Aneurysms of the pulmonary artery. *Circulation* 2015;131:310-6.
4. Stout KK, Daniels CJ, Aboulhosn JA, et al. 2018 AHA/ACC guideline for the management of adults with congenital heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation* 2019;139:e698-800.

---

**KEY WORDS** bicuspid pulmonary valve, congenital disease, giant aneurysm, pulmonary artery

---

**APPENDIX** For supplemental videos, please see the online version of this paper.